

“Underlying Health Conditions One”

Analyses of Underlying Health Conditions Among Minorities’ Children in Southwest Houston, Texas and The Roles Parents Play to Complicate the Inabilities to Control Them

¹Dr. Catherine W. Kisavi-Atatah, ²Dr. Park E. Atatah, ³Dr. Angela Branch-Vital, ⁴Dr. Donna Y. Laws, and ⁵Dr. Jasmine J. Opusunju

¹Ph.D. Assistant Professor, Prairie View A&M University, Prairie View, TX, USA

²Ph.D., Assistant Professor, Prairie View A&M University, Prairie View, TX, USA

³Ph.D. Associate Professor, Departmental Head, Prairie View A&M University, Prairie View, TX, USA

⁴Ed.D. Assistant Professor, Prairie View A&M University, Prairie View, TX, USA

⁵Dr PH. Clinical Assistant Professor, Baylor University, Waco, TX, USA

ABSTRACT

This quasi-experimental project’s study investigated the relationship between multidiscipline approaches in addressing the issues associated with minorities’ children between the ages of 6 and 18 weights, obesity, BMI, and their parental physical, mental, social, and overall parental involvements or lack of involvements in their children’s health wellbeing outcomes. The study questioned the outcomes by analyzing the population/patients, intervention/indicator, comparator/control, with no pediatric obesity educational program, outcomes such as impact the dietary habits, and time for over a period of 4 weeks in six months (PICOT). The project study employed the roles of parental activities, in activities, and possibly “out of control” parental supervisions as indicators. The project study used a singular “case study” to collect data statistics using Child Feeding Questionnaire (CFQ) survey instrument developed by Birch et al. (2007). The project’s study investigated 20 children/parents using ANOVA to analyze pretests versus posttests, and scorecards. The project’s study found that there were statistically significant differences between dependent and independent variables in pretests versus posttests in more than 30 plus tables and figures. The project’s study recommends that collective collaborations between physicians and multidiscipline healthcare practitioners is needed in addressing some of the resolvable issues associated with children’s weights, obesity, BMI, and blood sugar levels AIC was way to manage their underlying health conditions. The study recommends that a longer time study should be conducted; as to identify how long and how sustainable the parents and the children are

committed the process. This study recommends that future studies should limit their items of analysis to not more than 3 to 5 items per a project as to reduce data statistics outputs' overruns. The project's study concludes that if all the above itemized recommendations among others are carefully implemented, we should be able to maintain and possibly gain some sustainable positive social changes in the future to come.

KEYWORDS: *Child Feeding Questionnaire (CFQ), PICOT, BMI, Weight, Height, Obesity, Overeating, Blood Sugar Levels, A1C, Underlying Conditions, Out-of-Control Children, CORONAVIRUS, COVID-19, and Healthy 2020*

INTRODUCTION

The purpose of this quasi-experimental quantitative research study was to examine the relationships between children overeating habits and in activities with their weight gains and Body Mass Index (BMI) blood sugar A1C levels associated with minorities in Southwest Houston, Texas. This study investigated 15 associative items; but, only the first two items were analyzed in this study due to the overwhelming database statistics it generated after the initial analyses. However, additional items such as item 3 to 15 will be addressed in the follow-up studies in the future. This study collected some monthly samples from 20 volunteered participants. Since the issues associated with CORONAVIRUS or COVID-19 has become prevalence among minorities in general, this study intends to shed some lights on minorities between the ages of 6 to 18 years children and the roles their parents may play or not play to control their weight gains and BMI blood sugar A1C levels collectively.

LITERATURE REVIEWED

The American Heart Association (2017) approximates that one out of every ten children are either overweight or obese. According to the same report, the prevalence of childhood obesity has more than tripled over the last three decades, with the health problem surpassing drug and substance abuse to become the number one childhood health concern. The excess weight places the children at the peril for development of serious health problems like diabetes, heart disease and asthma. Being overweight/obese in childhood is associated with a risk of obesity-related co morbidities in adulthood, adverse psychological problems, huge societal expenses and premature morbidity and mortality (see Grigorakis et al., 2016; foster et al., 2017; Fonvig et al., 2015; Fidelix et al., 2015 for more).

Childhood obesity predisposes children to other serious chronic and acute health problems, hence limiting the children from reaching their full potentials in childhood and adulthood (Pulgarón, 2014). Childhood obesity has remained a communal health issue that continues to affect children in their right to relish a long healthy life

free of avertible diseases. According to the American Heart Association (2017), childhood obesity is the number one trepidation for every parent, having surpassed the issue of drug abuse and smoking as top public health problem for children (See Roy & Soleiman, 2014; Pulgarón, 2014; Ogden et al., 2014; Martin et al., 2018; Mendes et al., 2016; Damaso et al., 2013; &Mameli et al., 2017 and more).

Obesity among children and adolescents is described as a body condition in which the BMI figure exceeds 30. The most causes of the condition are identified as physical inactivity and unhealthy diets with high calorie content with the weight effects moderated by society and environment factors. Given that obesity is a health risk factor, there is a need to address its incidence with the focus on achieving and maintaining a healthy body weight. This is particularly true for children and adolescents since obesity first presents in this age group with the best management results achieved. Cochrane et al. (2017) notes that obesity management is important, but it is not only concerned with weight management with input from different professionals required to achieve the desire outcome in terms of healthy weight loss. It further adds that a multidisciplinary approach is necessary to achieve the best outcomes. Foster et al. (2017) similarly notes that the best outcomes are achieved by combining input from different stakeholders since the conditions has implications that exceed weight and could extend to dieting, family support, and exercising. Based on this awareness, the study recommends that family-centered approaches should be applied when addressing weight management needs of children and adolescents since they are easily influenced by their families.

Fonvig et al. (2015) also mentions that a multidisciplinary approach reduced BMI. The article further added that the approach reduced liver fat, muscle fat and visceral adipose tissue volumes although these results were only noted after running the program for 1 year. The implication is that it presents positive metabolic effects (Fonvig et al., 2015). The same sentiments are expressed by Martin et al. (2018), mentioning that although obesity management is about weight control, it also has cognitive implications for children and adolescents since a healthy weight will typically be associated with good performance in school. This means that in addition to the weight, other professionals would be required to address the cognitive aspects of care. Torti et al. (2017) make similar when calling for a multidisciplinary approach by indicating that obesity management is all about lifestyle intervention and should be matched with support from the school and community.

Carayol et al. (2017) presents a novel perspective by indicating that although the weight management strategies have achieved some success in obesity control, the success is limited and could be improved by including proteomic factors since the condition has molecular mechanisms. The implication is that molecular biologists should also be included in obesity care provision since they can evaluate and interpret proteomic

results. Swift et al. (2014) mention that effective weight management to address obesity must include exercise programs along with calorific programs. The implication is that the medical personnel who manage the exercise program must work in concert with the personnel who work the calorific program since the two are different medical disciplines. Ogden et al. (2014) faults the current obesity intervention measures by noting that they have been largely ineffective since surveillance data shows that obesity prevalence remains high. The article adds that there is a need to for a multidisciplinary approach with input from different stakeholders to reverse the high prevalence noted.

Bocca et al. (2012) validates the recommendations for a multidisciplinary approach in obesity management by reporting that multidisciplinary care produces better outcomes when compared to usual-care programs and if used for at least one year. Mameli et al. (2017) expresses some reservations in using multidisciplinary weight management in children. The article notes that although a multidisciplinary approach is more effective when the whole program is followed, it is similarly accompanied by high dropout rates even before discernible results could be produced. The implication is that the approach might have positive effects, but it is demanding thereby causing high dropout rates (Mameli et al., 2017). The reviewed journal articles show that obesity among children and adolescents is best managed through a multidisciplinary approach although there could be a need to address dropouts to improve outcomes and effectiveness.

PICOT

The picot question has been presented as: “Among parents of obese or overweight children or adolescent from 6 to 18-year-old in a primary care pediatric setting (P), how does the implementation of a multidisciplinary pediatric obesity educational program (I) versus no pediatric obesity educational program (C) impact the dietary habits (O) over a period 4 weeks in 6 months (T)?”

P - Population/ Patient = how does the implementation of a multidisciplinary pediatric obesity educational program

I - Intervention/ Indicator = versus no pediatric obesity educational program

C - Comparator/control = no pediatric obesity educational program

- O - Outcome = impact the dietary habits
- T – Time = over a period of 4 weeks measurements for 6 months from January 5, 2020 to June 30, 2020.

CONCEPTUAL/ THEORETICAL FRAMEWORK

The prevalence of obesity has risen dramatically in the USA, with the rate increasing more than threefold during the past three decades. The most alarming is that childhood obesity most often will continue into adulthood and the consequences of obesity include coronary artery disease, diabetes, adverse psychological problems, large societal costs and premature morbidity and mortality (Grigorakis et al., 2016). About one in every three American kids and youths is either obese or overweight. Berge et al. (2016) presents the result of a quantitative research study into the effects of community-based childhood obesity prevention programs. The study acknowledged that obesity intervention programs have had limited success, particularly those targeting children. This failure is postulated to be the result of an inability to engage parents in the program. As such, there is a need to understand how community engagement affects the results of intervention programs. This problem is important to medical personnel since it addresses an ongoing concern in the form of identifying effective obesity prevention programs. Identifying the more effective programs and their approaches can facilitate effective resource allocation so that emphasis remains on programs that are likely to produce positive results. In this respect, medical personnel could use the research results to recommend greater community engagement in childhood-obesity intervention programs as a more effective strategy.

Many studies have proven that obesity intervention programs targeted at children have largely failed. The failure has been attributed to an inability to engage the whole community that would have a positive additive effect on outcomes. Majority of children aged 5 to 12 years watch, on average, almost two-and-a-half hours of television daily and most of this is commercially sponsored. In addition, around half of them now eat their evening meal in front of the television which has been recognized as an effective medium for the sale of food products with the vast majority of foods advertised to children being high in fat, salt and/or sugar and low in fiber. Changing social attitudes to childhood may influence the development of children's food preferences. Empirical evidence suggests that food availability and accessibility, parental role modeling, television viewing and child-parent interactions around food are all likely to be important.

Environmental factors like lack of physical activity are major contributors to the obesity epidemic and children who are obese or overweight are more likely to have functional limitations. Physical activity has been proven to lower the risk of obesity, but future studies are needed to determine barriers to participation among these children in recreation and sporting activities. By exploring the role of social networks and obesity, it shows that the obesity epidemic is affected by the complex interaction between the environment, genetic factors, and human behavior (Hong, Coker-Bolt, Anderson, Lee, & Velozo, 2016). The knowledge gap justifies the research

by determining whether there is any justification for multidisciplinary obesity program and community engagement in obesity intervention programs targeted at children.

DESIGN OF THE STUDY

The recruitment procedure entailed the identification of the potential participants based on the inclusion and exclusion criteria. Recruitment fliers with the Principal Investigator (PI) contact information was placed at the prospective clinic site. The convenient samples of patients were voluntarily enrolled after receipt of signed consent from their parents or guardians. Families responded with qualified children were invited for assessment where weight and height and BMI, and their blood sugar A1C were measured. Computer-generated random numbers was used to allocate parent-child pairs to the multidisciplinary pediatric obesity program. Parents and guardians were provided an informed consent while children and adolescents completed an assent (Bonnie et al., 2016).

METHODOLOGY

The study applied a quantitative approach that entailed by collecting data using one-group pre-test/post-test quasi-experimental design. A sample size of 20 participants was recruited from among patients of Family Clinic in Southeast Houston, Texas. Just to protect the interests of the clinic and the participants, the specifications were not needed in this study. The inclusion criteria were obesity diagnosis for more than one year, and parental/ guardian consent. The exclusion criteria were mental incompetence, deafness, blindness, terminal illness with an inability for self-care, and an inability to actively participate in provider-led education programs. The participant's blood pressure, weight, waist circumference, BMI and height were measured. Participants/parents were assessed on obesity knowledge, nutritional lifestyle, and physical activity with Child Feeding Questionnaire (CFQ) developed by Birch et al. (2007). The Child Feeding Questionnaire (CFQ) is a self-report measure to assess parental beliefs, attitudes, and practices regarding child feeding, with a focus on obesity proneness in children. The Nurse Practitioner who is also the principal investigator and physician will coordinate the multidisciplinary team (Reichert-Anderson et al., 2015).

The principle investigator/smet with the participants and their parents/guardians obtain a signed consent and complete a socio-demographic survey and administer the pre-test obesity knowledge Questionnaire. The participants/parents/guardians attended one-time individual obesity education session based on the teaching outline for 1 hr. There was a break every 15 minutes during the session to alleviate the emotional stress because of the sensitivity of the topic. There was a re-evaluation in four weeks after the education classes where the

participants/parents/guardians were asked to complete a posttest CFQ Questionnaire. Participants and parent/guardian were assessed on dietary patterns to promote healthy dietary habits and lifestyle behavioral changes according to the American Academy of Pediatrics' guidelines. The lifestyle modification counseling included promoting nutritious diet and physical activities (Reichert-Anderson et al, 2015). Personalized nutritional counseling was provided to overweight participants while a low-calorie balanced personalized diet was provided to the obese participants that seeks to achieve the target of (decrease in Body Mass Index (BMI) and reduction in childhood obesity (Roya & Soleiman, 2014).

HYPOTHESES

This study hypothesized two major research alternative hypotheses as shown below;

Alternative Hypothesis 1

There is a relationship between children overeating, in-activities, and the overall outcomes of their weight gains and Body Mass Index (BMI) blood sugar A1C levels

Alternative Hypothesis 2

There is a relationship between lack of parental supports, and inactivates their children overeating, in-activities, and the overall outcomes of their weight gains and Body Mass Index (BMI) blood sugar A1C levels

Critical questions as to why children overeat were;

1. "How often do you eat because you are lonely, bored, nervous, or upset?"
2. "How often do you get out of bed at night, go into the kitchen, and finish the remains of some food?"

Null Hypotheses

This study hypothesized two major research Null hypotheses as shown below;

Null Hypothesis 1

There is no relationship between children overeating, in-activities, and the overall outcomes of their weight gains and Body Mass Index (BMI) blood sugar A1C levels

Null Hypothesis 2

There is no relationship between lack of parental supports, and inactivates their children overeating, in-activities, and the overall outcomes of their weight gains and Body Mass Index (BMI) blood sugar A1C levels

INTERVENTION AND DATA COLLECTION

Data Collection Methods

The data collection methodology applicable for this study was through a quantitative approach with a pre-test/post-test quasi-experimental design. Fliers were placed at the prospective clinic for voluntary sign up of participants with parental/guardian consent. They can also be referred by a confidential doctor in a private clinic in Southwest Houston, Texas based on their BMI percentiles at or exceeding the 95th percentile according to the CDC normative scale. Data was collected by measuring weight, height and BMI. The obesity knowledge Questionnaire with socio-demographic survey Questionnaire was used as multiple and open-ended style questions to fully elicit the nutritional tendencies and lifestyle living of the participants.

Data Analysis Plan

The PICOT statement seeks to determine if a multidisciplinary pediatric obesity educational program compared to a non-multidisciplinary pediatric obesity educational program will impact the dietary habits changes in participants over a period 4 weeks repeatedly for six months.

PICOT Statement Interpretation

Population: The population is obese children/ adolescents between the ages of 6 years and 18 years. Most of these patients attend a primary care clinic where their interventions are varied depending on their medical condition and the pediatrician's approach.

Intervention: Most of the current pediatric obesity management regimes are centered on nutrition. However, there is a need to develop patient-centered pediatric obesity educational program that involves a multidisciplinary team in order to benefit from the combination of several proven strategies.

Control: A multidisciplinary pediatric obesity educational program approach will be compared to no pediatric obesity educational program.

Outcome: The desired outcome will be a desired impact on dietary habits in the participants in a period of 4 weekly for six months

Time: The time for this research is 4 weekly for six months.

PROJECT ANALYSIS

Statistical Analyses

This project will use quantitative data analyses as outlined below.

Data Analyses Plan

First tests of treatment differences in demographic characteristics was performed using χ^2 tests for definite variables and t tests for incessant variables. After collection of the data from both groups, a simple linear regression analysis was used to establish if the multidisciplinary pediatric obesity educational program will impact the dietary habits over a period 4 weeks.

Analyses will be performed using the available data. All statistical analysis was performed using IBM Statistical Package for Social Sciences (SPSS) Version 25. In addition, follow-up visits were used as likely predictors. Every predictor was analyzed using the linear regression model where predictors with significant correlation with impact on the dietary habits were included in linear regression model. Quantitative survey data was analyzed using SPSS version 25, SPSS Inc, Chicago, IL, USA for cross-tabulation analysis to assess respondents' ability to make changes with goals of healthier eating (Collins, Warren, Neve & Stokes, 2016).

PROJECT FINDINGS/RESULTS OF THE STUDY

Pending the conclusion and the findings/results of the project

Classification of Collected Raw Data from the Survey Instrument

Races of Participants

Whites 2

Blacks /African Americans 4

Hispanic/Black Hispanic Americans 12

Asians 1

Pacific Americans 1

N=20

Genders of Participants

Female=18

Male=2

N=20

Demographics

To protect the interests and the privacies of the participants, personal demographic information such ages, names, date of birth (DOB), drivers’ licenses (DL), addresses, and residential zip codes were not needed or obtained in this project.

Pretests Results/Findings of the Study

Pretests

The following questions are about eating and physical activity. Please mark the box that corresponds to the frequency in which each situation takes place.

Table 1: Reasons for eating excessively always never occasionally frequently usually

How often do you eat because you are lonely, bored, nervous, or upset?	6	0	6	5	3	20
How often do you get out of bed at night, go into the kitchen, and finish the remains of some food?	5	4	5	3	3	20

Table 1. The reasons for eating excessively based on the two pinpointed items (see table 1 above and more).

Table 2: Implementation of health care interventions

For each activity that you, your spouse, or your child do 15 minutes or more, write down the number of times each of you perform the activity during an average week.

	YOU	YOUR SPOUSE	OTHER	CHILD
Walking	10	5	3	2
Running/Jogging	7	6	5	2
Playing tag	9	4	3	4
Dancing/Aerobic dance	8	6	5	1
Hiking/climbing	0	0	0	0
Baseball/softball/Football	7	4	4	5
Basketball/Volleyball	0	0	0	0
Jumping rope	8	6	4	2
Soccer	5	7	5	2
Skateboarding/skating	5	7	4	4
Swimming laps	5	8	2	5
Bicycling	7	7	3	3
Tennis/badminton	2	8	3	2
Other				

Table 2. Showed that the participants were followed monthly by collecting the weights as well as their BMI as to see if there were any improvements health improvements after implementing the above interventions (see table 2 above and more).

Descriptive Statistics

Table 3. Pre-Test 1

Statistics

	How often do you eat because you are lonely, bored, nervous, or upset	How often do you get out of bed at night, go into the kitchen, and finish the remains of some food?
N Valid	20	20
Missing	0	0
Mean	3.4500	3.0000
Std. Error of Mean	.27601	.31623
Median	3.5000	2.5000
Mode	3.00 ^a	2.00
Std. Deviation	1.23438	1.41421
Variance	1.524	2.000
Skewness	-.246	.372
Std. Error of Skewness	.512	.512
Kurtosis	-.949	-1.336
Std. Error of Kurtosis	.992	.992
Range	4.00	4.00
Minimum	1.00	1.00
Maximum	5.00	5.00
Sum	69.00	60.00

a. Multiple modes exist. The smallest value is shown

Table 3 showed the ANOVA of pre-tests which was divided into 8 items as classified above due to the overwhelming data statistics generated by SPSS version 25. Table 1 analyzed the means, median; standard deviation, variances, Std. Error of skewness of .512, and 0 missing data of 20 participants (see Table 3 above and more).

Frequency Table

Table 4. How often do you eat because you are lonely, bored, nervous, or upset

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Never	1	5.0	5.0	5.0
	Occasionally	4	20.0	20.0	25.0
	Frequently	5	25.0	25.0	50.0
	Usually	5	25.0	25.0	75.0
	Always	5	25.0	25.0	100.0
	Total	20	100.0	100.0	

Table 4. Showed the cumulative frequency of 5 items such as Never, Occasionally, Frequently, Usually, and Always with no missing data of 100% Valid and Cumulative Frequencies (see table 4 above and more).

Table 5. How often do you get out of bed at night, go into the kitchen, and finish the remains of some food?

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Never	2	10.0	10.0	10.0
	Occasionally	8	40.0	40.0	50.0
	Frequently	3	15.0	15.0	65.0
	Usually	2	10.0	10.0	75.0
	Always	5	25.0	25.0	100.0
	Total	20	100.0	100.0	

Table 5. Showed the cumulative frequency of 5 items such as Never, Occasionally, Frequently, Usually, and Always with no missing data of 100% Valid and Cumulative Frequencies (see table 5 above and more).

Figure 1.




Figure 1 showed the mean of 3.45, Std. Deviation of 1.234 and N=20 (see Figure 1 above and more).

Figure 2.




Figure 2 showed the mean of 3.00, Std. Deviation of 1.414 and N=20 (see Figure 2 above and more).

Pre-Test T-Test

Table 6. One-Sample Statistics

	N	Mean	Std. Deviation	Std. Error Mean
How often do you eat because you are lonely, bored, nervous, or upset	20	3.4500	1.23438	.27601
How often do you get out of bed at night, go into the kitchen, and finish the remains of some food?	20	3.0000	1.41421	.31623

Table 6 was Pre-Test T-Test with N=20 no missing numbers, the lowest mean was 1.23 while the highest mean was 3.45, and the highest Std. deviation was 1.41

Table 7. One-Sample Test

Test Value = 0

	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval Lower
How often do you eat because you are lonely, bored, nervous, or upset	12.499	19	.000	3.45000	2.8723
How often do you get out of bed at night, go into the kitchen, and finish the remains of some food?	9.487	19	.000	3.00000	2.3381

Statistics

Table 7 showed a One Sample Test Sig (2 tailed) test with Internal Confidence Differences with SPSS version 25 set at 95% or 0.05. From items 1 and 2 there were no significant differences of .000&.000, from items 1 and 2 between dependent and independent variables. As such, the project accepted the Alternative Hypotheses that there were relationships between parents/children inactivity, food intakes, and BMI and weight gain in 4 weeks after 6 months. The project rejected the Null Hypotheses that there were no relationships between parents/children’s activities, food intakes and BMI as well as weight gain. However, the study found that there was a perfect relationship between dependent and independent variables at .000 or 100% relationship (see table 7 above and more).

Posttests Results/Findings of the Study

Table 8: Posttests

		How often do you eat into the kitchen, and because you are lonely, bored, nervous, or upset	How often do you get out of bed at night, go finish the remains of some food?
N	Valid	20	20
	Missing	0	0
Mean		2.6500	1.8500
Std. Error of Mean		.28354	.25418
Median		2.5455 ^a	1.6250 ^a
Mode		2.00	1.00
Std. Deviation		1.26803	1.13671
Variance		1.608	1.292
Skewness		.400	1.518
Std. Error of Skewness		.512	.512
Kurtosis		-.696	2.023
Std. Error of Kurtosis		.992	.992

Range	4.00	4.00		
Minimum	1.00	1.00		
Maximum	5.00	5.00		

- a. Calculated from grouped data.
- b. Multiple modes exist. The smallest value is shown

Table 8 showed the ANOVA of Post-tests 1 which was divided into 8 items as classified above due to the overwhelming data statistics generated by SPSS version 25. Table 20 analyzed the means, median; standard deviation, variances, Std. Error of skewness of .512, and 0 missing data of 20 participants; however, there were significant differences between item 8 of the Pre-Tests and Post-Tests (see Table 8 above).

Table 9: Frequency Distribution

How often do you eat because you are lonely, bored, nervous, or upset

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Never	4	20.0	20.0	20.0
	Occasionally	6	30.0	30.0	50.0
	Frequently	5	25.0	25.0	75.0
	Usually	3	15.0	15.0	90.0
	Always	2	10.0	10.0	100.0
	Total	20	100.0	100.0	

Table 9. Showed the cumulative frequency of 5 items such as Never, Occasionally, Frequently, Usually, and Always with no missing data of 100% Valid and Cumulative Frequencies.

Table 10: How often do you get out of bed at night, go into the kitchen, and finish the remains of some food?

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Never	10	50.0	50.0	50.0

Occasionally	6	30.0	30.0	80.0
Frequently	2	10.0	10.0	90.0
Usually	1	5.0	5.0	95.0
Always	1	5.0	5.0	100.0
Total	20	100.0	100.0	

Table 10. Showed the cumulative frequency of 5 items such as Never, Occasionally, Frequently, Usually, and Always with no missing data of 100% Valid and Cumulative Frequencies.

Figure 3: Posttests Histogram Distribution 1




Figure 3 showed the mean of 2.65, Std. Deviation of 1.27 and N=20 (see Figure 3 above and more).

Table 4: Posttests Histogram Distribution 2




Figure 4 showed the mean of 1.85, Std. Deviation of 1.14 and N=20 (see Figure 4 above and more).

Table 11: Post-Test T-Test

One-Sample Statistics

	N	Mean	Std. Deviation	Std. Error Mean
How often do you eat because you are lonely, bored, nervous, or upset	20	2.6500	1.26803	.28354
How often do you get out of bed at night, go into the kitchen, and finish the remains of some food?	20	1.8500	1.13671	.25418

Table 11 was Post-Test T-Test with N=20 no missing numbers, the lowest mean was 1.85, the lowest Std. deviation was 1.14 while the highest mean was 2.65, and the highest Std. deviation was 1.27 (see table 11 above and more).

Table 12: One-Sample Test

	Test Value = 0			Mean Difference	95% Confidence Interval of the Difference	
	t	df	Sig. (2-tailed)		Lower	Upper
How often do you eat because you are lonely, bored, nervous, or upset	9.346	19	.000	2.65000	2.0565	3.2435
How often do you get out of bed at night, go into the kitchen, and finish the remains of some food?	7.278	19	.000	1.85000	1.3180	2.3820

Table 12 showed a One Sample Test Sig (2 tailed) test with Internal Confidence Differences with SPSS version 25 set at 95% or 0.05. From items 1 and 2 there were no significant differences .000, and .000, from items 1to 2 between dependent and independent variables. As such, the project accepted the Alternative Hypotheses that there were relationships between parents/children inactivity, food intakes, and BMI and weight gain in 4 weeks after 6 months. The project rejected the Null Hypotheses that there were no relationships between parents/children’s inactivity, food intakes, and BMI as well as weight gain (see table 12 above and more).

Interpretations of the Results/Findings of the Study

Based on the overwhelming data generated through the comprehensive analyses of the pretests versus posttests, the first five top tables and first five figures were interpreted as to answer this quasi experimental quantitative research study. However, since the ANOVA of table 1 was already analyzed, table was not part of the data interpretation of the study’s findings and results. The study found in **pretests** that 1 our 20 participants or 5% claimed that they never eat whenever they are lonely, bored, nervous, or upset. At the same time 4 out 20 or 20% claimed eat occasionally, and 5 out 20 or 25% claimed they eat frequently, usually, and always. The study found that the participants get out of bed at night, go into the kitchen, and finish the remains of some food at 2 out 20 or 10% never, 8 out 20 or 40% occasionally, 3 out of 20 or 10% frequently, 2 out of 20 or 10% always, and 5 out of 20 or 25% claimed always.

On a contrary hand, the study found in **posttests** that 4 out of 20 participants or 20% claimed that they never eat whenever they are lonely, bored, nervous, or upset. At the same time 6 out of 20 or 30% claimed eat occasionally, and 5 out of 20 or 25% claimed they eat frequently, 15% claimed usually, and 10 claimed always. The study found that the participants get out of bed at night, go into the kitchen, and finish the remains of some food at 10 out of 20 or 50% never, 6 out of 20 or 30% occasionally, 2 out of 20 or 10% frequently, 1 out of 20 or 5% always, and 1 out of 20 or 5% claimed always.

Interpretations of Findings/Results from 15 Minutes or More Physical Activities Scorecards

After completing the project's intervention, the participants submitted their scorecards in their individual, family, and other's collective 15 minutes or more physical activities that they participated in the past 4 weeks in six months as indicated in Appendices 1 below. The data statistics from the scorecards showed that 11 out of 20 or 55% walked alone; while 4 out of 20 or 20% walked with their spouses, 0 out of 20 or 0% with others, and 5 out of 20 or 25% walked with their children. As for running/jogging 6 out of 20 or 30% walked alone, 5 out of 20 or 25% with their spouses, 3 out of 20 or 15% with others, and 6 out of 20 or 30% with their children. More interestingly, as for playing tag, the project found that 6 out of 20 or 30% with their children, same as in dancing/aerobic dance, 30% with their children in basketball/softball/football with children and 30% in jumping rope.

Furthermore, 7 out of 20 or 35% played soccer with their children, 4 out of 20 or 20% in skateboard/skating and in tennis/badminton. These improvements remain consistent in swimming laps at 25% with their children and so in bicycling (see Appendices 1 below). The above data statistics indicate that there was a significant improvement in parents and children physical activities. However, **it should be noted that the ending BMI was not measured due to the short-term of the intervention process.** In fact, the visible changes of BMI take time to materialize; and 4 weeks for six months was not enough. Conclusively, it should also be noted that these overrides and concerns were strongly suggested in the suggestion and recommendations' section of this project; as to get a more comprehensive applicable outcomes about BMI implications due to relationships between inactivity and food intakes among children's weight issues. In fact, it should be noted that there were significant relationships at .000 and .000 or 100% between the dependent and independent variables in this study.

Answers to PICOT Project Study's Research Questions

The picot question has been presented as: "Among parents of obese or overweight children or adolescent from 6 to 18-year-old in a primary care pediatric setting (P), how does the implementation of

a multidisciplinary pediatric obesity educational program (I) versus no pediatric obesity educational program (C) impact the dietary habits (O) over a period 4 weeks in six months (T)?”

As to answer the PICOT project study’s question as pinpointed above and below;

P - Population/ Patient = how does the implementation of a multidisciplinary pediatric obesity educational program

I - Intervention/ Indicator = versus no pediatric obesity educational program

C - Comparator/control = no pediatric obesity educational program

- O - Outcome = impact the dietary habits
- T – Time = over a period of 4 weeks in six months.

The conclusion and discussion and the interpretations of the project study’s findings/results shed some fundamental lights and provided the investigative answers about the project’s study investigations (please, see interpretations of findings/results of pretests and posttests, scorecards’ findings/results, and conclusion and discussion).

IMPLICATION OF THE STUDY

This project showed several significance and implications to participants, researcher/s, healthcare practitioners and others in several ways:

- First, the findings/results of this project should assist multidiscipline health care practitioners about the proper and possibly the most effective ways to manage children’s weight and BMI generally.
- Secondly, this project’s findings/results should shed some lights to the parents about the importance of their (parents’) involvements in their children’s overall healthcare activities outcomes.
- Third, the parents should know that their involvements should be holistic and not sometimes or occasionally.
- Fourth, the parents should be able to identify the important items in the activities’ intervention survey instrument/s that could lead to better, progressive, continuous, and possibly the most positive outcomes.
- Significantly, the parents should know that their involvements do not start and end at home alone; but, in schools, food intakes, physical activities, mental activities, psychological activities, other related activities, and overwhelming parental involvements in their children’s healthcare wellbeing.

- Finally, these findings/results of this project should assist health care practitioners about how to effectively, efficiently, and even proficiently positively manage issues associated with children's weight and BMI in general.

This project's findings/results summed that if the above pinpointed significance/implications are carefully implemented, the issues associated with children's weight and BMI implications and complications should be systematically or even symmetrically eliminated with time.

LIMITATION OF THE PROJECT'S STUDY

This project showed several limitations which are pinpointed below:

- First, this project was conducted and concluded as a quasi-experimental quantitative research study which lacks randomization of participants' selections and activities.
- Quasi-experimental quantitative research studies operate on limited randomization which leads to the inability for findings/results generalization.
- This project was conducted as a singular case study based on the location of the project; as well as the originalities of the collections of the participants data statistics of the study.
- Case study poses yet another limitation due to its inability to be generalized outside the case study's location.
- This study was conducted by primary researcher who worked in the location, knew the multidiscipline team/s, and may know the participants personally, interpersonally, and professionally. These relationships may pose biases on the part of the research team/s in general.
- Furthermore, the scorecards results were self-reported which posed some issues of research validities and reliabilities for the results and findings.
- Finally, the scope of this study is limited due to the size of the participants; which yet poses another limitation for overwhelming generalization outside the scope.

Significance or Implications

This project showed several significance and implications to participants, researcher/s, healthcare practitioners and others in several ways:

- First, the findings/results of this project should assist multidiscipline health care practitioners about the proper and possibly the most effective ways to manage children's weight and BMI generally.
- Secondly, this project's findings/results should shed some lights to the parents about the importance of their (parents') involvements in their children's overall healthcare activities outcomes.
- Third, the parents should know that their involvements should be holistic and not sometimes or occasionally.
- Fourth, the parents should be able to identify the important items in the activities' intervention survey instrument/s that could lead to better, progressive, continuous, and possibly the most positive outcomes.
- Significantly, the parents should know that their involvements do not start and end at home alone; but, in schools, food intakes, physical activities, mental activities, psychological activities, other related activities, and overwhelming parental involvements in their children's healthcare wellbeing.
- Finally, these findings/results of this project should assist health care practitioners about how to effectively, efficiently, and even proficiently positively manage issues associated with children's weight and BMI in general.

This project's findings/results summed that if the above pinpointed significance/implications are carefully implemented, the issues associated with children's weight and BMI implications and complications should be systematically or even symmetrically eliminated with time.

CONCLUSION AND DISCUSSION

As to understand the conclusion and discussion of this project, it is fair to note the primary purpose of the study. The project was to investigate if the adaptation of multidisciplinary approach would improve obesity and overweight management outcomes when compared to care provided solely by a primary care provider. This approach was based on a need to present a more efficient management approach that would guarantee the best outcomes. This need was galvanized by the fact that obesity and overweight conditions as a source of concern within the healthcare industry affecting emotional, social and physical health wellbeing. This project found that involvements of multidisciplinary was more effective in the improvements of obesity and overweight management outcomes when compared to care provided solely by a primary care provider.

The study noted and summed that the involvements of the parents of their children's activities and food intakes showed a more significant outcomes in the management of weight and BMI levels. The

study also concludes that there were overwhelming significant statistical differences in all areas especially in the visual pinpointed tables and figures in the importance and the effects of intervention in children's weight and BMI controllable management. As such, the study concludes that involving parents in their children's activities such as physical activities, discussions about the importance of food intakes at home, stress issues, isolations at home and schools, and other places, are fundamentally significant across the board. The data statistics children's/parents' physical activities indicate that there was a significant improvement in parents and children physical activities in the past 4 weeks. However, it should be noted that the ending BMI was not measured due to the short-term of the intervention implementation as shown in Appendix 1.

Above all, parents were advised that by demonstrating the effective, efficient, and proficient ways to achieve these pinpointed outcomes, due to parental holistic involvements in their children's activities is holistically more fundamental in achieving some positive social changes in managing children's weight and BMI physically, socially, mentally, physiologically and psychologically wellbeing in the final analyses. In conclusion, from a public health care policy standpoint, and even from a public policy standpoint, this project should unequivocally assist health care multidiscipline, the most accurate, effective, efficient, and proficient ways in addressing managing weight and IBM issues among children in general. Since the recent pandemic we currently faced in the US, evidence upon evidence have shown the underlying health conditions such as overweight, BMI, high blood sugar level A1C, in activities are corresponding critical factors in killing minorities whenever they are infested with **CORONAVIRUS also known as COVID-19**. This study shed some lights about how to effectively, efficiently, and proficiently control these underlying poor health conditions as to make majority of minorities who participated in this study healthier than ever; in the future to come. Hopefully, this study will possibly make some pinpointed participants new acquired eating and exercises' habits engagements as a part of their new **Healthy 2020 lifestyle**.

RECOMMENDATIONS (MAINTAINING/SUSTAINING THE CHANGE)

This project recommends the followings as to be able to maintain and sustain the noted changes based on its findings/results.

- An experimental research study needs to be conducted instead of quasi-experimental research study which poses limitations due its inability to apply its findings/results in other settings.

- Multiple case studies need to be conducted instead of singular case study based on its limited data originalities.
- Collective collaborations between physicians and multidiscipline healthcare practitioners need to be united on how to proactively address some of the resolvable issues associated with children’s weight and BMI management applications.
- The study recommends that a longer time study should be conducted; as to identify how long and how sustainable the parents and the children are committed the process.
- This study recommends that future studies should limit their items of analysis to not more than 3 to 5 items per a project as to reduce data outputs’ overruns. And finally,
- The researcher and possibly other researchers should make some of the data available to other health care practitioners for findings/results implementations.

In summary, the project concludes that the above itemized recommendations should be able to maintain and possibly sustain some positive social changes for some long time to come; for all majority of minorities’ children in Southwest Houston, Texas and possibly beyond.

ACKNOWLEDGEMENTS

We want to use this opportunity to thank all the scholars for their contributions in completing this complex and complicated social scientific research study which underscored underlying health conditions in the US which are detrimental to minorities in general when dealing with CORONAVIRUS or COVID-19. We want to specifically thank the “**Confidential Clinic**” in Southwest Houston, Texas for allowing us to use their facility in completing this critical children/parents’ related project’s research study.

CONFLICT OF INTERESTS

We share no conflict of interests in this study because it was self-supported and funded.

REFERENCES

- 1 American Heart Association. (2017). *Overweight in children*. Retrieved from http://www.heart.org/HEARTORG/HealthyLiving/HealthyKids/ChildhoodObesity/OverweineChildren_UCM_304054_Article.jsp

- 2 Birch, L., Fisher, J., Grimm-Thomas, K., Markey, C., Sawyer, R. and Johnson, S. (2018). *Confirmatory factor analysis of the Child Feeding Questionnaire: a measure of parental attitudes, beliefs and practices about child feeding and obesity proneness.*
- 3 Bonnie J, Ronsen R, Roth S, Luke C, Nida S, Coombs L et al. (2016). The Fit Family Challenge: A Primary Care Childhood Obesity Pilot Intervention. *Journal of the American Board of Family Medicine.* 29(4), 434-443.
- 4 Bocca, G., Corpeleijin, E., Stolk, R. & Sauer, P. (2012). Results of a multidisciplinary treatment program in 3-year-old to 5-year-old overweight or obese children: a randomized controlled clinical trial. *Arch Pediatr Adolesc Med.*, 166(12), 1109-1115. doi:10.1001/archpediatrics.2012.1638
- 5 Berge, J. M., Jin, S., Hanson-Bradley, C., Doty, J., Jagaraj, K., Braaten, K. & Doherty, W. J. (2016). Play it Forward! A Community-Based Participatory Research Approach to Childhood Obesity Prevention. *Fam Syst Health*, 34(1), 15-30. doi: 10.1037/fsh0000116
- 6 Carayol, J., Chabert, C., Di Cara, A., Armensie, C., Vlasesia, A. & Haper. J. (2017). Protein quantitative trait locus study in obesity during weight-loss identifies a leptin regulator. *Nature Communications* 8, 2084 (2017). Retrieved from <https://www.nature.com/articles/s41467-017-02182-z#Abs1>
- 7 Cochrane, A., Dick, B., King, N., Hills, A. & Kavanagh, D. (2017). Developing dimensions for a multicomponent multidisciplinary approach to obesity management. *BMC Public Health*, 17, 814. Retrieved from <https://doi.org/10.1186/s12889-017-4834-2>
- 8 Collins, C. E., Warren, J., Neve. M., & Stokes, B. J. (2016). Measuring Effectiveness of Dietetic Interventions in Child Obesity. *Article Review.* Available at file:///C:/Users/owner/AppData/Local/Temp/pr60002_906_922.pdf
- 9 Damaso, A., de Piano, A., Campos, R., Corgosinho, F., Siegfried, W. ... & Tock, L. (2013). Multidisciplinary approach to the treatment of obese adolescents: effects on cardiovascular risk factors, inflammatory profile, and neuroendocrine regulation of energy balance. *International Journal of Endocrinology*, Volume 2013, Article ID 541032. doi: 10.1155/2013/541032. Retrieved from <https://www.hindawi.com/journals/ije/2013/541032/>
- 10 Fidelix, Y., de Farias, J., Lofrano-Prado, M., Guerra, R., Cardel, M. & do Prado, W. (2015). Multidisciplinary intervention in obese adolescents. *Einstein (Sao Paulo)*, 13(3). doi: 10.1590/S1679-45082015AO3339. Retrieved from http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1679-45082015000300388

- 11 Fonvig, C., Chabanova, E., Ohrt, J., Nielsen, L., Pedersen, O. ... & Holm, J. (2015). Multidisciplinary care of obese children and adolescents for one year reduces ectopic fat content in liver and skeletal muscle. *BMC Pediatrics*, 15, 196. doi: 10.1186/s12887-015-0513-6
- 12 Foster, C., Moore, J., Singletary, C. & Skelton, J. (2017). Clinical obesity: physical activity and family-based obesity treatment. *Wiley Online Library*, 8(1), 68-79. Retrieved from <http://onlinelibrary.wiley.com/doi/10.1111/cob.12230/full#publication-history>
- 13 Grigorakis, D., Georgoulis, M., Psarra, G., Tambalis, K., Panagiotakos, D., & Sidossis, L. (2016). Prevalence and lifestyle determinants of central obesity in children. *European Journal of Nutrition*, 55(5), 1923-1931. doi:10.1007/s00394-015-1008-9
- 14 Hong, I., Coker-Bolt, P., Anderson, K. R., Lee, D., & Velozo, C. A. (2016). Relationship Between Physical Activity and Overweight and Obesity in Children: Findings From the 2012
- 15 Karnik, S., & Kanekar, A. (2014). Childhood obesity: a global public health crisis. *International Journal of Preventive Medicine*, 3(1), 1.
- 16 Mameli, C., Krakauer, J., Krakauer, N., Bosetti, A., Ferrari, C. ... & Zuccotti, G. (2017). Effects of a multidisciplinary weight loss intervention on overweight and obese children and adolescents: 11 years of experience. *PLoS One*, 12(7), e0181095. doi: 10.1371/journal.pone.0181095
- 17 Mendes, A., Ieker, A., De Castro, T., Avelar, A. & Nardor, N. (2016). Multidisciplinary programs for obesity treatment in Brazil: A systematic review. *Revista de Nutricao*, 29(6). doi: 10.1590/1678-98652016000600011. Retrieved from http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1415-527320160006000867
- 18 Martin, A., Booth, J., Laird, Y., Sproule, J., Reilly, J. & Saunders, D. (2018). Physical activity, diet and other behavioural interventions for improving cognition and school achievement in children and adolescents with obesity or overweight. 1-185. *Cochrane Database of Systematic Reviews*. Retrieved from
- 19 Ogden, C., Carroll, M., Kit, B. & Flegal, K. (2014). Prevalence of childhood and adult obesity in the United States. *Jama Network Journals*, 311(8), 806-814. Retrieved from <https://jamanetwork.com/journals/jama/fullarticle/1832542>
- 20 Pulgarón, E. R. (2014). Childhood obesity: A review of increased risk for physical and psychological co-morbidities. *Clinical Therapeutics*, 35(1), A18-A32.
- 21 Roy K & Soleiman F. (2014). Controlling childhood obesity: A systematic review on strategies and challenges. *J Res Med Sci*. 19(10): 993–1008.

- 22 Reichert-Anderson P, Yang P, Yang C, Jacobson M & Bernstein H. (2015). A Multidisciplinary Approach to Childhood Obesity Prevention in Pediatric Primary Care. *Journal of the Academy of Nutritionists and Dieticians*. 115(9). <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD009728.pub3/full>
- 23 Swift, D., Johannsen, N., Lavie, C., Earnest, C. & Church, T. (2014). The role of exercise and physical activity in weight loss and maintenance. *ELSEVIER Publication*, 56(4), 441-447. Retrieved from: [http://www.online.pcd.com/article/S0033-0620\(13\)00165-5/fulltext](http://www.online.pcd.com/article/S0033-0620(13)00165-5/fulltext)
- 24 Torti, J., Liug, T., Borowitz, M., Johnson, J. A., Sharma, A. & Cambell-Shere, D. (2017). The 5As team patient study: patient perspectives on the role of primary care in obesity management. *BMC Family Practice*, 18(19). Retrieved from <https://link.springer.com/article/10.1186/s12875-017-0596-2>
- 25 Zolotarjova, J., ten Velde, G. & Vreugdenhil, A. (2018). Effects of multidisciplinary interventions on weight loss and health outcomes in children and adolescents with morbid obesity. *Obesity Reviews*, 19(7), 931-946. doi: 10.1111/obr.12

APPENDICES

Appendices 1

15 Minutes or More Physical Activities with/between Parents/Children

For each activity that you, your spouse, or your child do 15 minutes or more, write down the number of times each of you perform the activity during an average week.

	YOU	YOUR SPOUSE	OTHER	CHILD
Walking	11	4	0	5
Running/Jogging	6	5	3	6
Playing tag	7	2	0	6
Dancing/Aerobic dance	8	2	4	6
Hiking/climbing	0	0	0	0
Baseball/softball/Football	7	2	2	9
Basketball/Volleyball	0	0	0	6
Jumping rope	6	2	4	6
Soccer	5	3	5	7
Skateboarding/skating	2	3	2	4
Swimming laps	3	3	4	5
Bicycling	7	3	2	5
Tennis/badminton	2	2	7	4
Other				