

## PANCREATIC CANCER IN MID- REGION OF LIBYA; RETROSPECTIVE STUDY FROM JULY 2005 TO JAN 2015

**RAMADAN ELAMYAL\* & NAJI ZUBIA\*\***

\*MD Consultant Surgeon, Department of Surgery, Misurata Cancer Center, Misurata, Libya.

\*\*Senior Registrar, Department of Surgery, Misurata Cancer Center, Misurata, Libya.

### **ABSTRACT**

**INTRODUCTION:** *the pancreas is a glandular organ of dual composition with both exocrine and endocrine components. Tumors of exocrine pancreases are rare with more than 95% being malignant. METHODS & MATERIALS:* *this is a retrospective clinico-histopathological study of 89 cases registered in our center from July 2005 to Jan 2015. The details of patients were retrieved from patients' files. Information retrieved included socio-demographic data, clinical presentation, anatomical site, gross appearance, tumor stage, histopathological type and grade, presence of metastasis. RESULTS:* *during the study period, a total of 89 malignancies were registered. Of these, 48 (53.9%) were histopathologically confirmed pancreatic cancer. The diagnosis in the remaining 41 patients was based on clinical and radiological basis. Male to female ratio of 1.6:1. The age ranged from 35 to 90 years with a median age of 63 years. Fifty seven patients (64%) presented with abdominal pain, 41 patients (46%) were jaundiced. The head was the most frequent site for pancreatic tumor in 48 (54%) cases followed by tail, body. Adenocarcinoma was the most common histopathological tumor in 43(90%) patients. Thirty (33.7%) patients were presented in locally advanced stages and 54(60.6 %) patients had metastasis. CONCLUSION:* *Our study was limited by being a retrospective, a relatively small number of patients and representing the experience of a single cancer center so, larger scale prospective studies are recommended to study the clinicodemographic character of this disease in the country.*

**KEY WORDS:** *Pancreatic Cancer, clinical, demographic, retrospective.*

### **INTRODUCTION:**

The pancreas is a glandular organ of dual composition with both exocrine and endocrine components [1, 2]. Tumors of exocrine pancreases are rare with more than 95% being malignant [3]. Several risk factors for pancreatic

cancer have been identified, such as smoking [4], age [5], family history [6], and diabetes [7]. The majority of pancreatic cancers are of exocrine origin and majority of adenocarcinomas type, commonly involving the head and body of the pancreas [8]. Cancer of pancreas is most commonly seen in the 7th and 8th decades with a median age of 71 years at diagnosis. Early pancreatic cancer has no specific symptoms and signs and there is no established screening strategy to detect it early. However, at the time of diagnosis, 24% of patients have locally advanced disease and 54% have distant metastases [9]. The purpose of the study is to study exocrine pancreatic cancers clinico-pathologically to know the various types, grades and clinical manifestations.

## **METHODS AND MATERIALS:**

This is a retrospective clinico-histopathological study of 89 cases registered in our Center from July 2005 to Jan 2015. The details of patients were retrieved from patients' files. Information retrieved included socio-demographic data, clinical presentation, anatomical site, gross appearance, tumor stage, histopathological type and grade, presence of metastasis.

## **RESULTS:**

- During the study period, a total of 89 malignancies were registered. Of these, 48 (53.9%) were histopathologically confirmed pancreatic cancer. The diagnosis in the remaining 41 patients was based on clinical and radiological basis. The number of males was 55 (62%) and the number of females was 34 (38%) with a male to female ratio of 1.6:1. The age ranged from 35 to 90 years with a median age of 63 years. Table 1.

**Table 1 Demographic data of 135 patients with colorectal Cancer.**

<b>Gender</b>	
Male	55 (62%)
Female	34 (38%)
<b>Mean age (SD), range</b>	63

- Fifty seven patients (64%) presented with abdominal pain, 41 patients (46%) were jaundiced, 29 patients (32%).Thirty seven patients (42%) had history of diabetes and 41 patients (46%) are smokers. Table (2).

**Table 2 Distribution of patients according to clinical presentation**

Symptoms & Signs	No (%)
Abdominal pain	57 (64%)
Obstructive Jaundice	41 (46%)
Loss of weight	29 (32%)
Anorexia	23 (26%)
Nausea& Vomiting	21 (23%)
Malaise	07 (7%)
Dyspnea& Cough	05 (5%)
Fever	02 (2%)
Subcutaneous nodules	01
PMH Diabetes	(1%)
History of smoking	37(42%)
	41(46%)

- The head was the most frequent site for pancreatic tumor in 48 (54%) cases followed by tail, body and diffuse in 10 (11%), 8 (9%) and 23(25%) cases, respect. Table 3.

**Table3 show site distribution of pancreatic cancer.**

Site of tumors	No (%)
Head	48(54%)
Tail	10(11%)
Body	8 (9%)
More than one subsite(Diffuse)	23(25%)

- Microscopically, adenocarcinoma was the most common histopathological type in 43(90%) patients. Carcinoid, Mucinous cystadenocarcinoma and acinar cell carcinomas accounted for 2 (4%), 1 (2%) and 1(2%) patients, respectively. Only one patient (2%) was mixed adenocarcinoma and carcinoid. Table 4.

**Table 4: show histopathology types**

Histopathology*	No (%)
<b>Adenocarcinoma</b>	<b>43 (90%)</b>
<b>Carcinoid</b>	<b>2 (4%)</b>
<b>Mucinous cyst adenocarcinoma</b>	<b>1 (2%)</b>
<b>Acinar cell carcinoma</b>	<b>1 (2%)</b>
<b>Mixed adenocarcinoma &amp; carcinoid</b>	

\*Histopathology was confirmed in 48 patients.

- Regarding extent of disease, only 5 (5.6%) patients were identified as being in early stages, 30 (33.7%) patients were presented in locally advanced stages and 54(60.6%) patients had metastasis. Table 5.

**Table 5: show extent of the disease.**

Extent of disease	No (%)
<b>Early stage</b>	<b>5 (5.6%)</b>
<b>Locally advanced</b>	<b>30(33.7%)</b>
<b>Metastasis</b>	<b>54(60.7%)</b>

**DISCUSSION:**

The two main demographic risk factors for pancreatic cancer are advancing age and male gender[10]. In the present study, the median age was 63 years and majority of cases of adenocarcinomas were found in the fifth and seventh decade (41.7% each). Compared to the median age of 72 years in United State [11], our patients present at approximately 9 years younger median age. This may be attributed to overall younger population pool in Libya or may be due to different biology of the cancer. Dohert et al [12] showed that seventy five (75%) of pancreatic cancer

patients presented with weight loss, obstructive jaundice and upper abdominal pain. Upper abdominal pain (53%) and weight loss (40%) were also common first symptoms [13]. In our study, sixty-four percent (64%) of pancreatic cancer patient presented with abdominal pain, 46% with obstructive jaundice and 32% with loss of weight. Other modes of presentations in our study were anorexia (23 patients 26%), nausea & vomiting (21 patients 25%), Malaise (7 patients 7%), Dyspnea & cough (5 patients 5%), fever (2 patients 2%) and subcutaneous nodules (one patient, 1%). No patients in our study were presented with bone pain. Diabetes mellitus and cigarette smoking may be possible risk factors for development of carcinoma of exocrine pancreas [14]. In this study there was 37 cases (42%) had diabetes mellitus and 41 cases (46%) were smokers. The majority of pancreatic cancers are adenocarcinomas. These tumours originate in the epithelial cells lining the pancreatic duct, form gland-like structures, and account for 90% of all pancreatic cancers [1,15,16]. In the present study, adenocarcinoma constituted the majority (90% patients) and is comparable to the study by Kumar NV [17] in which (89.7%) of cases were adenocarcinomas. Almost one third of our patients present at advanced stage and sixty percent present with distant metastasis compared to 53% in United State of America [18]. This may be due to the scarcity of specialized centers and investigations facilities so, a considerable number of patients remain undiagnosed or diagnosed at the advanced stage of the disease.

## CONCLUSION:

The study was limited by its retrospective nature, a relatively smaller number of patients and representing the experience of a single cancer center so, larger scale prospective studies are recommended to study the clinical and demographic character of this disease in the country.

## REFERENCES:

1. Rosai J. Pancreas and ampullary region. In: Rosai and Ackerman's Surgical Pathology. 10th ed. Missouri: Elsevier Mosby; 2011.
2. Hollingsworth MA. Proteins expressed by pancreatic duct cells and their relatives. *Ann N Y Acad Sci* 1999; 880:38-49.
3. Hruban RH, Donahue DL. The Pancreas. In: Kumar, Abbas, Fasuto, Aster, editors. Robbins and Cotran Pathologic Basis of Disease. 8th ed. Philadelphia: Elsevier Saunders; 2010. p. 891-903.
4. Warshaw AL, Castillo CF. Pancreatic carcinoma. *N Engl J Med* 1992; 326:455-65.
5. Ghadirian P, Lynch HT, Krewski D. Epidemiology of pancreatic cancer: an overview. *Cancer Detect Prev* 2003; 27: 87-93.
6. Lynch HT, Smyrk T, Kern SE, *et al.* Familial pancreatic cancer: a review. *Semin Oncol* 1996; 23:251-75.

7. Everhart J, Wright D. Diabetes mellitus as a risk factor for pancreatic cancer: a meta-analysis. *JAMA* 1995; 273:1605-9.
8. Dhir V, Mohandas K. Epidemiology of digestive tract cancers in India IV. Gall bladder and pancreas. *Indian J Gastroenterol* 1999; 18(1): 24-8.
9. Cancer Facts & Figures. American Cancer Society. Available at <http://www.cancer.org/Research/CancerFactsFigures/index>. Accessed: October 8, 2014.
10. Gold E.B., and Goldin B. Epidemiology of and risk factors for pancreatic cancer. *SurgOncolClin North Am* 1998; 7:67-91.
11. Howlader N, Noone AM, Krapcho M, Neyman N, AminouR, Waldron W, et al, editors. SEER Cancer Statistics Review 1975-2008 National Cancer Institute SEER Cancer Statistics Review 1975-2008 National Cancer Institute. Bethesda (MD):National Cancer Institute; 2011.
12. Doherty GM, Way LW. Pancreas. In: Way LW, Doherty GM, editors. Current surgical diagnosis and treatment. New York:Lange Medical Books; 2003. pp. 602–12.
13. Sonnefeld T, Nyberg B, Perbeck L. The effect of palliative biliodigestive operations for unresectable pancreatic carcinoma. *J Acta Chir Scand Suppl.* 1986; 530:47–50.
14. Shi C, Hruban RH, Klein AP. Familial pancreatic cancer. *Arch Pathol Lab Med* 2009; 133: 365–74.
15. Cowgill SM, Muscarella P. The genetics of pancreatic cancer. *Am J Surg* 2003; 186: 279-86.
16. Li D, Jiao L. Molecular epidemiology of pancreatic cancer. *Int J Gastrointest Cancer* 2003; 33:3-14.
17. Kumar NV. Clinical study of carcinoma pancreas [dissertation]. Mangalore: Rajiv Gandhi Univ.; 2007.
18. American Cancer Society. Cancer Facts & Figures 2011. USA:American Cancer Society, Inc.: 2011. p. 1-55.