


North Asian International Research Journal Consortium

*North Asian International Research Journal
Of
Multidisciplinary*

Chief Editor

Dr. Nisar Hussain Malik



NAIRJC JOURNAL PUBLICATION

North Asian
International
Research Journal Consortium



Welcome to NAIRJC

ISSN NO: 2454 - 2326

North Asian International Research Journal is a multidisciplinary research journal, published monthly in English, Hindi, Urdu all research papers submitted to the journal will be double-blind peer reviewed referred by members of the editorial board. Readers will include investigator in Universities, Research Institutes Government and Industry with research interest in the general subjects

Editorial Board

J.Anil Kumar Head Geography University of Thirvanathpuram	Sanjuket Das Head Economics Samplpur University	Adgaonkar Ganesh Dept. of Commerce, B.S.A.U Aruganbad
Kiran Mishra Dept. of English,Ranchi University, Jharkhand	Somanath Reddy Dept. of Social Work, Gulbarga University.	Rajpal Choudhary Dept. Govt. Engg. College Bikaner Rajasthan
R.D. Sharma Head Commerce & Management Jammu University	R.P. Pandday Head Education Dr. C.V.Raman University	Moinuddin Khan Dept. of Botany SinghaniyaUniversity Rajasthan.
Manish Mishra Dept. of Engg, United College Ald.UPTU Lucknow	K.M Bhandarkar Praful Patel College of Education, Gondia	Dr. Ashish Kumar Department of Management Studies L.N.Mithila University Darbhanga , Bihar
Tihar Pandit Dept. of Environmental Science, University of Kashmir.	Simnani Dept. of Political Science, Govt. Degree College Pulwama, University of Kashmir.	Ashok D. Wagh Head PG. Dept. of Accountancy, B.N.N.College, Bhiwandi, Thane, Maharashtra.
Neelam Yadav Head Exam. Mat.K..M .Patel College Thakurli (E), Thane, Maharashtra	Nisar Hussain Dept. of Medicine A.I. Medical College (U.P) Kanpur University	M.C.P. Singh Head Information Technology Dr C.V. Rama University
Ashak Husssain Head Pol-Science G.B, PG College Ald. Kanpur University	Khagendra Nath Sethi Head Dept. of History Sambalpur University.	Rama Singh Dept. of Political Science A.K.D College, Ald.University of Allahabad

Address: -North Asian International Research Journal Consortium (NAIRJC) 221 Gangoo, Pulwama, Jammu and Kashmir, India -192301, Cell: 09086405302, 09906662570, Ph. No: 01933-212815, Email: nairjc5@gmail.com, info@nairjc.com Website: www.nairjc.com



Existence and stability of Collinear libration point L_1 in the Restricted Three Body problem when the Primaries are Triaxial Rigid Bodies and Source of Radiations, Perturbation Effects Act in Coriolis and Centrifugal forces.

B. K. MANDAL*

*Department of Mathematics, Govt. Women's Polytechnic, Bokaro, Jharkhand, India

ABSTRACT:-

This paper deals with the stationary solutions of the collinear libration point L_1 of the planar restricted three body problem when the primaries are triaxial rigid bodies and source of radiations, perturbation effects act in coriolis and centrifugal forces with one of the axes as the axis of symmetry and its equatorial plane coinciding with the plane of motion. It is seen that there are five libration points two triangular and three collinear. It is further observed that the collinear points are unstable, while the triangular points are stable for the mass parameter $0 \leq \mu < \mu_{crit}$ (the critical mass parameter). It is further seen that the triangular points have long or short periodic elliptical orbits in the same range of μ .

Key words: Restricted Three Body Problem; Libration Point; Rigid Body; Source of Radiations; Coriolis force; Centrifugal force.

1. INTRODUCTION:

It is well known that the classical planar restricted three body problem possesses five libration points, two triangular and three collinear. The collinear libration points L_1 , L_2 , L_3 are unstable, while the two equilateral libration points L_4 , L_5 are stable for $\mu < \mu_{crit} = 0.0385208965 \dots \dots \dots$ Szebehely. Winter showed that the stability of the two equilateral points is due to the existence of coriolis terms in the equations of motion written in a synodic co-ordinate system.

In recent times many perturbing forces i.e., oblateness and radiation forces of the primaries, coriolis and centrifugal forces, variation of the masses of the primaries and of the infinitesimal mass etc., have been included in the study of the restricted three body problem. In the case of restricted three body problem where both the primaries are oblate spheroids whose equatorial plane coincides with the plane of motion, the location of libration points and their stability in the Liapunov sense has been studied by Vidyakin. For the case, where the bigger primary is an oblate spheroid whose equatorial plane coincides with the plane of motion, Subba Rao and Sharma



have studied the stability of libration points. A similar problems has been studied by El-Shaboury. Khanna and Bhatnagar have studied the problem when the smaller primary is a triaxial rigid body.

Sharma, Taqvi and Bhatnagar have studied the problem when the bigger primary is a triaxial rigid body as well as source of radiation.

In this paper, we consider the primaries are triaxial rigid bodies and source of radiation, perturbation effects act in coriolis and centrifugal forces with one of the axes as the axis of symmetry and their equatorial plane coinciding with the plane of motion. Further we assume that the primaries are moving without rotation about their centre of mass in circular orbits. An attempt is made to study the existence and location of libration point L_1 .

Mandal (2017) found collinear libration point L_1 in 2+2 body problem when perturbation effects act in coriolis and centrifugal forces, small primary is a radiating body and bigger primary is a triaxial rigid body.

2. EQUATIONS OF MOTION

We shall adopt the notation and terminology of Szebehely. As a consequence, the distance between the primaries does not change and is taken equal to one; the sum of the masses of the primaries is also taken as one. The unit of time is so chosen as to make the gravitational constant unity. Using dimensionless variables, the equations of motion of the infinitesimal mass m_3 in a synodic co-ordinate system (x, y) are

$$\ddot{x} - 2n \dot{y} = \frac{\partial \Omega}{\partial x}$$

and

$$\ddot{y} - 2n \dot{x} = \frac{\partial \Omega}{\partial y}, \quad (1)$$

$$\text{where } \Omega = \sum_{i=1}^2 \left[\frac{1}{2} n^2 \mu_i r_i^2 + (1 - \mu_i) \left\{ \frac{\mu_i}{r_i} + \frac{\mu_i}{2m_i r_i^3} (I_{1i} + I_{2i} + I_{3i} - 3I_i) \right\} \right] \quad (2)$$

McCusky

$$\left. \begin{aligned} \mu_1 &= 1 - \mu, \mu_2 = \mu \\ r_i^2 &= (x - x_i)^2 + y^2, \quad (i = 1, 2) \\ x_1 &= \mu, x_2 = -1 + \mu \end{aligned} \right\} \quad (3)$$

$$p_1 = \frac{\text{Radiation pressure due to bigger primary}}{\text{Gravitation force due to bigger primary}} \ll 1.$$

$$p_2 = \frac{\text{Radiation pressure due to smaller primary}}{\text{Gravitation force due to smaller primary}} \ll 1.$$

Here, we have applied effect of small perturbation in coriolis and centrifugal forces with the help of perturbations α and β the unperturbed value of both being unity

$$\alpha = 1 + \epsilon, |\epsilon| \ll 1$$

$$\beta = 1 + \epsilon', |\epsilon'| \ll 1$$

Hence Ω change to Ω' the equilibrium points are obtained from equations

$$\Omega'x = \Omega'y = 0.$$

Here μ is the ratio of the mass of the smaller primary to the total mass of the primaries and $0 < \mu \leq \frac{1}{2}$.

That is, $\mu = \frac{m_2}{m_1+m_2} \leq \frac{1}{2}$ with $m_1 \geq m_2$ being the masses of the primaries.

I_{1i}, I_{2i}, I_{3i} ($i = 1, 2$) are the principal moments of inertia of the triaxial rigid body of mass m_i ($i = 1, 2$) at its centre of mass, with a_i, b_i, c_i ($i = 1, 2$) as lengths of its semi-axes. I_i ($i = 1, 2$) is the moment of inertia about a line joining the centre of the rigid body of mass m_i ($i = 1, 2$) and the infinitesimal body of mass m_3 and is given by

$$I_i = I_{1i}l_{1i}^2 + I_{2i}m_{1i}^2 + I_{3i}n_{1i}^2 \quad (i = 1, 2).$$

where l_{1i}, m_{1i}, n_{1i} ($i = 1, 2$) are the direction cosines of the line with respect to its principal axes.

Here, we have also assumed that the principal axes of m_1 and m_2 are parallel to the synodic axes $O(xyz)$.

The axes $O(xyz)$ have been defined by Szebehely.

The mean motion, n , is given by

$$n^2 = 1 + \sum_{i=1}^2 \frac{3}{2} (2A_{1i} - A_{2i} - A_{3i}), \quad (4)$$

where $A_{1i} = \frac{a_i^2}{5R^2}, A_{2i} = \frac{b_i^2}{5R^2}, A_{3i} = \frac{c_i^2}{5R^2}, (i = 1, 2)$

and R is the distance between the primaries.

Here we are neglecting the perturbation in the potential between m_1 and m_2 due to radiation pressure because m_1 and m_2 are supposed to be sufficiently large.

Ω in the eq. (2) can also be written as

$$\Omega = \sum_{i=1}^2 \left[\frac{1}{2} n^2 \mu_i r_i^2 + \frac{\mu_i}{r_i} + \frac{\mu_i}{2r_i^3} (2\sigma_{1i} - \sigma_{2i}) - \frac{3\mu_i}{2r_i^5} (\sigma_{1i} - \sigma_{2i}) y^2 - P_i \frac{\mu_i}{r_i} \right]$$

where $\sigma_{1i} = A_{1i} - A_{3i}$ and $\sigma_{2i} = A_{2i} - A_{3i} (i = 1, 2)$.

We assume that σ_{1i} and $\sigma_{2i} \ll 1 (i = 1, 2)$.

The mean motion gives in the eq. (4), becomes

$$n^2 = 1 + \sum_{i=1}^2 \frac{3}{2} (2\sigma_{1i} - \sigma_{2i}) \quad (5)$$

It may be noted that the mean motion, n , is independent of the solar radiation pressure $p_i (i = 1, 2)$.

3. LOCATION OF LIBRATION POINT L_1

Equations (1) permits an integral analogous to Jacobi integral

$$\dot{x}^2 + \dot{y}^2 - 2\Omega + C = 0$$

The libration points are the singularities of the manifold

$$F(x, y, \dot{x}, \dot{y}) = \dot{x}^2 + \dot{y}^2 - 2\Omega + C = 0$$

Therefore, these points are the solutions of the equations

$$\Omega_x = 0, \Omega_y = 0$$

For “ L_1 ” coefficient of x

$$\begin{aligned} L_1 &= x \left[(1 + \epsilon') \left(1 + \frac{3}{2} A_1 \right) \left\{ \mu - \frac{1}{2} + \frac{1}{8\mu} (4 - \mu) \sigma_1 - \frac{1}{8\mu} (4 + 3\mu) \sigma_2 \right\} \right. \\ &\quad - (1 - \mu) \left\{ -\frac{1}{2} + \frac{1}{8\mu} (4 - \mu) \sigma_1 - \frac{1}{8\mu} (4 + 3\mu) \sigma_2 \right\} \times \left\{ 1 + \frac{33}{8} \sigma_1 - \frac{33}{8} \sigma_2 + \epsilon' \right\} \\ &\quad - \mu \left\{ \frac{1}{2} + \frac{1}{8\mu} (4 - \mu) \sigma_1 - \frac{1}{8\mu} (4 + 3\mu) \sigma_2 \right\} \times \left\{ 1 + \epsilon' - \frac{3}{2\mu} \sigma_1 (1 - 3\mu) - \frac{3}{2\mu} \sigma_2 (-1 + 2\mu) \right\} \\ &\quad \left. - \frac{3}{2} (1 - \mu) A_1 \times -\frac{1}{2} + \frac{45}{16} (1 - \mu) A_2 \right] \\ &= x \left[\sigma_1 \left(\frac{33}{16} + \frac{3}{4} - \frac{45}{16} + 3\mu - \frac{33}{16} \mu - \frac{9}{4} \mu - \frac{3}{2} \mu + \frac{45}{16} \mu \right) + \sigma_2 \left(-\frac{33}{16} - \frac{3}{4} + \frac{45}{16} + \frac{33}{16} \mu + \frac{3}{4} \mu - \frac{45}{16} \mu \right) \right] \\ &= x [\sigma_1 \times 0 + \sigma_2 \times 0] = 0 \end{aligned}$$

Coefficient of “ y ”

$$\begin{aligned} y &\left[(1 + \epsilon') \left(\frac{1}{2} + \frac{3}{4} A_1 \right) \left\{ \sqrt{3} - \frac{4\sqrt{3}}{9} \epsilon' + \frac{\sqrt{3}}{12\mu} (4 - 23\mu) \sigma_1 + \frac{\sqrt{3}}{12\mu} (-4 + 19\mu) \sigma_2 \right\} \right. \\ &\quad - (1 - \mu) \frac{\sqrt{3}}{2} \left\{ 1 - \frac{4}{9} \epsilon' + \frac{1}{12\mu} (4 - 23\mu) \sigma_1 + \frac{1}{12\mu} (-4 + 19\mu) \sigma_2 \right\} \times \left\{ 1 + \frac{33}{8} \sigma_1 - \frac{33}{8} \sigma_2 + \epsilon' \right\} \\ &\quad - \mu \times \frac{\sqrt{3}}{2} \left\{ 1 - \frac{4}{9} \epsilon' + \frac{1}{12\mu} (4 - 23\mu) \sigma_1 + \frac{1}{12\mu} (-4 + 19\mu) \sigma_2 \right\} \\ &\quad \times \left\{ 1 + \epsilon' - \frac{3}{2\mu} \sigma_1 (1 - 3\mu) - \frac{3}{2\mu} \sigma_2 (-1 + 2\mu) \right\} - \frac{3\sqrt{3}}{4} (1 - \mu) A_1 + \frac{3\sqrt{3}}{2} (1 - \mu) A_2 \\ &\quad \left. - \frac{45\sqrt{3}}{16} (1 - \mu) A_2 \right] \end{aligned}$$

$$\begin{aligned}
 &= y \left[\sigma_1 \left(-\frac{33\sqrt{3}}{16} + \frac{3\sqrt{3}}{4} - \frac{3\sqrt{3}}{2} + \frac{45\sqrt{3}}{16} + \frac{33\sqrt{3}}{16}\mu - \frac{9\sqrt{3}}{4}\mu + \frac{3\sqrt{3}}{2}\mu + \frac{3\sqrt{3}}{2}\mu - \frac{45\sqrt{3}}{16}\mu \right) \right. \\
 &\quad \left. + \sigma_2 \left(\frac{33\sqrt{3}}{16} - \frac{3\sqrt{3}}{4} + \frac{3\sqrt{3}}{2} - \frac{45\sqrt{3}}{16} - \frac{33\sqrt{3}}{16}\mu + \frac{3\sqrt{3}}{2}\mu - \frac{3\sqrt{3}}{4}\mu - \frac{3\sqrt{3}}{2}\mu + \frac{45\sqrt{3}}{16}\mu \right) \right] \\
 &= y[\sigma_1 \times 0 + \sigma_2 \times 0] = 0
 \end{aligned}$$

Coefficient of \dot{x}

$$\begin{aligned}
 &n\alpha \left[\dot{x} \left\{ -\frac{\sqrt{3}}{2} + \frac{2\sqrt{3}}{9}\epsilon' - \frac{\sqrt{3}}{24\mu}(4 - 23\mu)\sigma_1 - \frac{\sqrt{3}}{24\mu}(-4 + 19\mu)\sigma_2 \right\} \right] \\
 &= \left\{ 1 + \frac{3}{4}(2\sigma_1 - \sigma_2) + \epsilon \right\} \left[\dot{x} \left\{ -\frac{\sqrt{3}}{2} + \frac{2\sqrt{3}}{9}\epsilon' - \frac{\sqrt{3}}{6\mu}\sigma_1 + \frac{23\sqrt{3}}{24}\sigma_1 + \frac{\sqrt{3}}{6\mu}\sigma_2 - \frac{19\sqrt{3}}{24}\sigma_2 \right\} \right] \\
 &= \dot{x} \left[-\frac{\sqrt{3}}{2} + \frac{2\sqrt{3}}{9}\epsilon' - \frac{\sqrt{3}}{2}\epsilon + \sigma_1 \left(\frac{5\sqrt{3}}{24} - \frac{\sqrt{3}}{6\mu} \right) + \sigma_2 \left(-\frac{5\sqrt{3}}{12} + \frac{\sqrt{3}}{6\mu} \right) \right]
 \end{aligned}$$

Coefficient of \dot{y}

$$\begin{aligned}
 &n\alpha \left[\dot{y} \left\{ \mu - \frac{1}{2} + \frac{1}{8\mu}(4 - \mu)\sigma_1 - \frac{1}{8\mu}(4 + 3\mu)\sigma_2 \right\} \right] \\
 &= \dot{y} \left[-\frac{1}{2} + \mu + \epsilon \left(\mu - \frac{1}{2} \right) + \sigma_1 \left(-\frac{1}{8} - \frac{3}{4} + \frac{3}{2}\mu + \frac{1}{2\mu} \right) + \sigma_2 \left(-\frac{3}{4}\mu - \frac{1}{2\mu} \right) \right] \\
 &= \dot{y} \left[-\frac{1}{2} + \mu + \epsilon \left(\mu - \frac{1}{2} \right) + \sigma_1 \left(-\frac{7}{8} + \frac{3\mu}{2} + \frac{1}{2\mu} \right) + \sigma_2 \left(-\frac{3}{4}\mu - \frac{1}{2\mu} \right) \right]
 \end{aligned}$$

4. STABILITY OF COLLINEAR LIBRATION POINT L_1

First we consider the point lying in $(\mu - 2, \mu - 1)$.

For this point, $r_2 < 1$, $r_1 > 1$, we have

$$\Omega'_{x \circ y} = \frac{3\sqrt{3}}{2} \left[\mu - \frac{1}{2} + \frac{11}{18}\epsilon' + \frac{\sigma_1}{24\mu}(8 - 47\mu + 89\mu^2) + \frac{\sigma_2}{24\mu}(-8 + 9\mu - 37\mu^2) \right]$$

$$\Omega'_{x \circ x} = \frac{3}{4} + \frac{5}{4}\epsilon' + \frac{3}{16\mu}\sigma_1(15\mu^2 - 8 + 19\mu) + \frac{3}{16\mu}\sigma_2(-31\mu^2 + 8 - \mu) > 0$$

$$\Omega' y^{\circ} y = \frac{9}{4} + \frac{7}{4} \epsilon' + \frac{3}{16\mu} \sigma_1 (8 + 29\mu - 15\mu^2) + \frac{3}{16\mu} \sigma_2 (-8 - 7\mu + 15\mu^2) < 0$$

Similarly, for the points lying in $(\mu-1, 0)$ and $(\mu, \mu+1)$

$$\Omega' x^{\circ} y = 0, \Omega' x^{\circ} x > 0 \text{ and } \Omega' y^{\circ} y < 0$$

$$\text{Because } \Omega' x^{\circ} x \Omega' y^{\circ} y - (\Omega' x^{\circ} y)^2 < 0$$

The discriminant is positive and the four roots of the characteristic equation can be written as $\lambda_1 = s, \lambda_2 = -s, \lambda_3 = it$ and $\lambda_4 = -it$ (s and t are real) so the motion around the collinear points is unbounded and consequently the collinear points are unstable.

5. CONCLUSION:

In the restricted three body problems, when the primaries are triaxial rigid bodies as well as source of radiations, perturbation effects act in coriolis and centrifugal forces. There are five libration points, three collinear and two triangular. The collinear points are unstable for all values of the mass parameter μ . Such that $0 < \mu \leq \frac{1}{2}$ and the triangular points are stable for $\mu < \mu_{\text{crit}}$.

REFERENCES:

1. P.V.Subba Rao and R.K.Sharma, Astron, Astrophys, 43 (1975), 381
2. M.Khanna and K.B.Bhatnagar, Indian J.Pure appl. math 29(10), 1998.
3. R.K. Sharma, Z.A. Taquvi and K.B.Bhatnagar. Indian J.Pure appl. math 32(2) 255-266 February 2001.
4. D.N.Garain, B.K.Mandal and S.N.Adhikary (2007), Triangular equilibrium points and stability in the restricted three bodies when bigger primary is a triaxial rigid body and perturbation effects act in coriolis and centrifugal forces ARJPS 10 (1-2) 2007, pp, 27-31
5. B.K.Mandal (2017) IJRDT vollume-7, issue 4 April 2017 pp. 4-10.



Publish Research Article

Dear Sir/Mam,

We invite unpublished Research Paper, Summary of Research Project, Theses, Books and Book Review for publication.

Address:- North Asian International Research Journal Consortium (NAIRJC)
221, Gangoo Pulwama - 192301

Jammu & Kashmir, India

Cell: 09086405302, 09906662570,

Ph No: 01933212815

Email: nairjc5@gmail.com, info@nairjc.com

Website: www.nairjc.com

